

19



Europäisches Patentamt
European Patent Office
Office européen des brevets



11 Publication number:

0 496 989 A1

12

EUROPEAN PATENT APPLICATION

21 Application number: **91121536.6**

51 Int. Cl.⁵: **A01N 43/50, //(A01N43/50,
33:18,33:20,25:30,25:02)**

22 Date of filing: **01.01.92**

30 Priority: **31.01.91 US 648083**

43 Date of publication of application:
05.08.92 Bulletin 92/32

84 Designated Contracting States:
**AT BE CH DE DK ES FR GB GR IT LI LU NL PT
SE**

71 Applicant: **AMERICAN CYANAMID COMPANY**
1937 West Main Street P.O. Box 60
Stamford Connecticut 06904-0060(US)

72 Inventor: **Ong, Chungjian Jerry**
26 Glenview Drive
Warren, New Jersey 07059(US)

74 Representative: **Wächtershäuser, Günter, Dr.**
Tal 29
W-8000 München 2(DE)

54 **Herbicide emulsifiable concentrate compositions of imidazolinone herbicides.**

57 The present invention relates to a herbicide emulsifiable concentrate composition which comprises 0.5% to 5% by weight of an imidazolinyl acid; 0% to about 40% by weight of a 2,6-dinitroaniline derivative; 2% to 6% of an alkylarylsulfonic acid; ethylene oxide/propylene oxide block copolymer and alkphenol polyethylene oxide condensate emulsifying agents; an antigellation agent; and an aromatic solvent.

EP 0 496 989 A1

BACKGROUND OF THE INVENTION

United States Patent 4,749,404 describes herbicidal liquid concentrate compositions of salts of 2-(4-isopropyl-4-methyl-5-oxo-2-imidazolin-2-yl)-3-quinolinecarboxylic acid which may be combined with an organic dinitroaniline composition and United States Patent 4,822,405 describes concentrated oil in water emulsion combination compositions of water soluble salts of imidazolinone herbicides and dinitroaniline herbicides. However, the ratio of herbicides which may be employed is limited because of problems with crystallization which may occur with more concentrated formulations especially when storing the compositions at low temperatures. In addition, these compositions employ salts of imidazolinones and require both aqueous and organic solvents.

SUMMARY OF THE INVENTION


The present invention relates to a herbicidal emulsifiable concentrate composition which comprises 0.5% to 5% by weight of an imidazolinyl acid; 0% to about 40% by weight of a 2,6-dinitroaniline derivative; 2% to 6% of an alkylarylsulfonic acid; ethylene oxide/propylene oxide block copolymer and alkylphenol polyethylene oxide condensate emulsifying agents; an antigelation agent; and an aromatic solvent.

It is therefore an object of the present invention to advantageously provide an emulsifiable concentrate combination composition in which all of the active herbicidal agents are dissolved in a single solvent which may be stored at lower temperatures and for a longer period of time than suspension concentrate compositions or concentrated oil in water emulsion compositions.

It is also an object of the present invention to provide herbicidal emulsifiable concentrate compositions of free acids of imidazolinone herbicides and 2,6-dinitroaniline herbicides which are physically and chemically stable.

DETAILED DESCRIPTION OF THE INVENTION


A preferred embodiment of the invention is herbicidal emulsifiable concentrate compositions comprising on a weight basis about 0.5% to 5% of an imidazolinyl acid represented by structural formula I




(I)

wherein

- W is oxygen or sulfur;
 R₁ is C₁-C₄ alkyl;
 R₂ is C₁-C₄ alkyl or C₃-C₆ cycloalkyl; and when R₁ and R₂ are taken together with the carbon to which they are attached they may represent C₃-C₆ cycloalkyl optionally substituted with methyl;
 X is hydrogen, halogen, hydroxyl or methyl;
 Y and Z are independently hydrogen, halogen, C₁-C₆ alkyl, C₁-C₄ alkoxyalkyl, hydroxyloweralkyl, C₁-C₆ alkoxy, C₁-C₄ alkylthio, phenoxy, C₁-C₄ haloalkyl, nitro, cyano, C₁-C₄ alkylamino, diloweralkylamino, or C₁-C₄ alkylsulfonyl group or phenyl optionally substituted with one C₁-C₄ alkyl, C₁-C₄ alkoxy or halogen; and when taken together, Y and Z may form a ring in which YZ are represented by the structure: -(CH₂)_n-, where n is an integer of 3 or 4, provided that X is hydrogen; or



where L, M, Q and R are independently hydrogen, halogen, C₁-C₄ alkyl or C₁-C₄ alkoxy;
0% to about 40% of a herbicidal 2,6-dinitroaniline derivative represented by structural formula II



(II)

wherein

R₃ is hydrogen, C₂-C₄ alkyl or chloroethyl;

R₄ is C₂-C₅ alkyl, chloroethyl, 2-methylallyl or cyclopropylmethyl;

R₅ is hydrogen, methyl or amino;

R₆ is trifluoromethyl, C₁-C₃ alkyl, -SO₂NH₂ or SO₂CH₃;

2% to 6% of an alkylarylsulfonic acid; 2% to 6% of an ethylene oxide/propylene oxide block copolymer; 1% to 6% of an alkylphenol polyethylene oxide condensate; 0.5% to 10% of an antigellation agent and a sufficient amount of an aromatic solvent to total 100%.

A most preferred embodiment of the present invention comprises on a weight basis 0.5% to 5% of a formula I imidazolinyl acid, 20% to 40% of a formula II 2,6-dinitroaniline derivative, 2% to 6% of an alkylarylsulfonic acid, 2% to 6% of an ethylene oxide/propylene oxide block copolymer, 1% to 6% of an alkylphenol polyethylene oxide condensate, 0.5% to 10% of an antigellation agent and a sufficient amount of an aromatic solvent to total 100%.

Surprisingly, it has been found that the compositions of the present invention provide physically and chemically stable herbicidal emulsifiable concentrate compositions of formula (I) imidazolinyl acids and formula (II) 2,6-dinitroaniline derivatives which remain free-flowing and homogeneous for extended periods of time, and also remain physically and chemically stable after repeated freezing and thawing cycles without precipitating insoluble solids.

The alkylarylsulfonic acid of the present invention is an especially important element of the present compositions and should be present in at least an amount equal to about 1.2 times the percent by weight of the imidazolinyl acid to ensure that the acid completely dissolves in the aromatic solvent. Alkylarylsulfonic acids suitable for use in the compositions of the invention include C₈-C₁₈ alkylbenzenesulfonic acids, with dodecylbenzenesulfonic acid being most preferred.

Antigellation agents are used in the compositions of the present invention to ensure that the compositions remain homogeneous and free-flowing. Antigellation agents which are suitable for use in the compositions of the present invention include N-methylpyrrolidone, cyclohexanone, alcohols such as ethanol and methanol, glycols such as propylene glycol and ethylene glycol and the like with N-methylpyrrolidone being most preferred.

A preferred group of ethylene oxide/propylene oxide block copolymers suitable for use in the compositions of the present invention are butyl-omega-hydroxypoly(oxypropylene)block polymer with poly(oxyethylene) having an average molecular weight in a range of 2,400 to 3,500, with alpha-butyl-omega-hydroxy-ethylene oxide-propylene oxide block copolymers having an HLB of 12 and a viscosity at 25°C of 2,000 CPS, (TOXIMUL® 8320, Stepan Chemical Co.) being a most preferred member of this class of emulsifiers.

Preferred alkylphenol polyethylene oxide condensates suitable for use in the compositions of the

present invention are the nonylphenol ethoxylates, with nonylphenol ethoxylate (9 to 10 mols of ethylene oxide)(FLO MO®9N, DeSoto, Inc., Sellers Chemical Div.) being a most preferred member of this class of emulsifiers.

Aromatic solvents suitable for use in the compositions of this invention include aromatic hydrocarbon and chlorinated aromatic hydrocarbon solvents and mixtures thereof; such as toluene, xylenes, polynuclear aromatic hydrocarbons such as naphthalenes and alkylnaphthalenes and mixtures thereof, many of which are available from the fractionation of crude oil and in general have distillation ranges in the temperature range of from about 140°C to 305°C, with those having a distillation range of from about 183° to 280°C being most preferred, and are commercially available under a variety of tradenames; and mono or polychlorobenzene and toluenes.

Imidazoliny acids particularly useful in the compositions of the present invention are 2-(4-isopropyl-4-methyl-5-oxo-2-imidazolin-2-yl)-3-quinoline-carboxylic acid, 5-ethyl-2-(4-isopropyl-4-methyl-5-oxo-2-imidazolin-2-yl)nicotinic acid, 2-(4-isopropyl-4-methyl-5-oxo-2-imidazolin-2-yl)nicotinic acid and 2-(4-isopropyl-4-methyl-5-oxo-2-imidazolin-2-yl)-5-(methoxymethyl)nicotinic acid.

2,6-Dinitroaniline derivatives which are especially suitable for use in the compositions of the present invention are pendimethalin and trifluralin.

The emulsifiable concentrate compositions of the present invention may conveniently be prepared by admixing the formula (I) imidazoliny acid with a solution of the desired antigellation agent, alkylarylsulfonic acid, ethylene oxide/propylene oxide block copolymer, alkylphenol polyethylene oxide condensate and aromatic solvent until all of the solids are dissolved, then the formula (II) 2,6-dinitroaniline derivative is added to the solution and mixing is continued until a homogeneous solution is obtained.

The following examples are provided to further illustrate the compositions of the present invention but are not limitative of the invention described herein.

EXAMPLE 1

Preparation of herbicidal emulsifiable concentrate compositions

Solid 5-ethyl-2-(4-isopropyl-4-methyl-5-oxo-2-imidazolin-2-yl)nicotinic acid (2.62g, 93.4% pure, 2.62% on a weight basis) is added to a stirred solution of N-methylpyrrolidone (4g, 4% on a weight basis) and dodecylbenzenesulfonic acid (4g, 4% on a weight basis). The mixture is stirred until all of the solids dissolve. An aromatic hydrocarbon mixture (C₁₀ to C₁₃) aromatics having a distillation range of from about 226°-279°C (AROMATIC® 200, Exxon)(45.68g, 45.68% on a weight basis) is added to the mixture and stirring is continued until a homogeneous solution of obtained. Pendimethalin (35.7g, 92.3% pure, 35.7% on a weight basis) is added to the homogeneous solution and stirring is continued until the solids dissolve. Next, nonylphenol ethoxylate (9 to 10 mols of ethylene oxide)(4g, 4% on a weight basis) and alpha-butyl-omega-hydroxy-ethylene oxide propylene oxide block copolymer having a HLB of 12 and a viscosity at 25°C of 2,000 CPS (4g, 4% on a weight basis) are added to the mixture and stirring is continued until a homogeneous solution is obtained which is then filtered through 1.5 micron filter paper to obtain a clear dark brown solution having a density at 20°C of 1.0683 g/mL.

Following the procedure of Example 1, but using the compounds listed in Table I below gives the emulsifiable concentrate compositions listed in Table II below.

TABLE I

Herbicidal Imidazolinyl Acid

- a. 5-ethyl-2-(4-isopropyl-4-methyl-5-oxo-2-imidazolin-2-yl)nicotinic acid

Herbicidal 2,6-Dinitroaniline Derivative

- b. pendimethalin
- c. trifluralin

Alkylarylsulfonic Acid

- d. dodecylbenzene sulfonic acid

Antigelation Agent

- e. N-methylpyrrolidone
- f. cyclohexanone
- g. dipropylene glycol

Ethylene Oxide/propylene Oxide Block Copolymer

- h. alpha-butyl-omega-hydroxy-ethylene oxide-propylene oxide block copolymer having an HLB of 12 and a viscosity at 25°C of 2,000 CPS

Alkylphenol Polyethylene Oxide Condensate

- i. nonylphenol ethoxylate (6 mols of ethylene oxide)
- j. nonylphenol ethoxylate (9-10 mols of ethylene oxide)
- k. nonylphenol ethoxylate (11 mols of ethylene oxide)

- 5
- l. nonylphenol ethoxylate (13 mols of ethylene oxide)
 - m. nonylphenol ethoxylate (15 mols of ethylene oxide)
 - 10 n. nonylphenol ethoxylate (30 mols of ethylene oxide)

Aromatic Solvent

- 15
- o. aromatic hydrocarbon mixture (C_{10} to C_{13} aromatics, distillation range 226° - 279° C) (AROMATIC®200, Exxon)
 - 20 p. aromatic hydrocarbon mixture (C_9 to C_{12} aromatics, distillation range 183° - 210° C) (AROMATIC®150, Exxon)
- 25
- 30
- 35
- 40
- 45
- 50
- 55

TABLE II
Emulsifiable Concentrate Compositions

Compo- sition	Imidazo- linyl acid/ % W/W	2,6-Dinitro- aniline/ % W/W	Alkylaryl- sulfonic acid/ % W/W	Anti- gelation agent/ % W/W	Ethylene oxide/ propylene oxide block copolymer/ % W/W			Alkylphenol polyethylene oxide Conden- sate/% W/W	Aromatic Solvent/ % W/W
1	a/2.62	b/35.70	d/4.0	e/4.0	h/4.0	h/4.0	j/4.0	j/4.0	o/45.68
2	a/2.40	b/33.40	d/4.0	e/8.0	h/4.0	h/4.0	j/4.0	j/4.0	o/44.20
3	a/2.60	b/35.80	d/4.0	e/8.0	h/4.0	h/4.0	j/2.0	j/2.0	o/43.60
4	a/2.40	b/33.40	d/4.0	e/4.0	h/4.0	h/4.0	j/4.0	j/4.0	o/48.20
5	a/2.40	b/33.40	d/4.0	e/4.0	h/3.0	h/3.0	j/4.0	j/4.0	o/49.20
6	a/2.40	b/33.40	d/4.0	e/4.0	h/2.0	h/2.0	j/4.0	j/4.0	o/50.20
7	a/2.40	b/33.40	d/4.0	e/4.0	h/3.0	h/3.0	j/3.0	j/3.0	o/50.20
8	a/2.40	b/33.40	d/4.0	e/4.0	h/2.0	h/2.0	j/3.0	j/3.0	o/51.20
9	a/2.60	b/35.80	d/4.0	e/0.5	h/4.0	h/4.0	j/4.0	j/4.0	o/50.90
10	a/2.60	b/35.80	d/4.0	f/1.0	h/4.0	h/4.0	j/4.0	j/4.0	o/48.60
11	a/2.60	b/35.80	d/4.0	g/1.0	h/4.0	h/4.0	j/4.0	j/4.0	o/48.60
12	a/2.60	b/35.80	d/4.0	e/8.0	h/4.0	h/4.0	n/1.0	n/1.0	o/44.60
13	a/2.50	b/34.60	d/4.0	e/1.0	h/4.0	h/4.0	n/1.0	n/1.0	o/51.90
14	a/2.65	c/29.66	d/3.0	e/2.0	h/3.0	h/3.0	j/3.0	j/3.0	p/56.69

TABLE II (continued)

Compo- sition	Imidazo- linyl acid/ % W/W	2,6-Dinitro- aniline/ % W/W	Alkylaryl- sulfonic acid/ % W/W	Anti- gelation agent/ % W/W	Ethylene oxide/ propylene oxide block copolymer/ % W/W		Aromatic Solvent/ % W/W
					oxide Conden- sate/% W/W	polyethylene oxide	
15	a/2.65	c/29.66	d/3.0	e/4.0	h/3.0	j/3.0	p/54.69
16	a/2.45	c/29.66	d/3.0	e/4.0	h/3.0	i/3.0	p/54.89
17	a/2.45	c/29.66	d/3.0	e/4.0	h/3.0	k/3.0	p/54.89
18	a/2.45	c/29.66	d/3.0	e/4.0	h/3.0	l/3.0	p/54.89
19	a/2.45	c/29.66	d/3.0	e/4.0	h/3.0	m/3.0	p/54.89
20	a/2.62	c/29.66	d/4.0	e/8.0	h/4.0	j/4.0	p/47.72
21	a/2.45	c/29.44	d/3.0	e/4.0	h/3.0	j/3.0	o/55.11

EXAMPLE 2**Low temperature stability of emulsifiable concentrate compositions of the invention**

Two 10mL samples of compositions 2 and 4 prepared in the above example are placed in a constant temperature bath maintained at 5°C and the temperature of the samples is allowed to equilibrate for two


hours. Seeds of pendimethalin crystals and 5-ethyl-2-(4-isopropyl-4-methyl-5-oxo-2-imidazolin-2-yl)nicotinic acid powder are added to the samples. After 48 hours, the samples are visually inspected for evidence of crystallization. The temperature of the bath is then lowered in 1°C increments holding the sample at each temperature for 48 hours until crystallization is observed. A comparative control composition containing on a weight basis 31.71% pendimethalin, 2.26% 5-ethyl-2-(4-isopropyl-4-methyl-5-oxo-2-imidazolin-2-yl)nicotinic acid, 0.5% ammonium hydroxide, 27.42% aromatic mixture (AROMATIC® 200, Exxon), 6.04% surfactants and 32.07% water is also evaluated similarly except the initial testing temperature is set at 15°C instead of 5°C. The results of these experiments are summarized in Table III below.

TABLE III**Low Temperature Experiments**

<u>Composition</u>	<u>Crystallization Temperature °C</u>
Control	13°C
2	-1°C
4	-3°C
9	0°C
11	0°C

Claims

1. A herbicidal emulsifiable concentrate composition comprising about 0.5% to 5% by weight of an imidazoliny acid; 0% to about 40% by weight of a 2,6-dinitroaniline derivative; 2% to 6% by weight of an alkylarylsulfonic acid; ethylene oxide/propylene oxide block copolymer and alkylphenol polyethylene oxide condensate emulsifying agents; an antigelation agent; and an aromatic solvent.
2. The composition according to Claim 1 wherein the imidazoliny acid has the structural formula (I) below

**(I)**

wherein

W is oxygen or sulfur;


R₁ is C₁-C₄ alkyl;

R₂ is C₁-C₄ alkyl or C₃-C₆ cycloalkyl; and when R₁ and R₂ are taken together with the carbon to which they are attached they may represent C₃-C₆ cycloalkyl optionally substituted with methyl;


X is hydrogen, halogen, hydroxyl or methyl;

Y and Z are independently hydrogen, halogen, C₁-C₆ alkyl, C₁-C₄ alkoxyalkyl, hydroxyloweralkyl, C₁-C₆ alkoxy, C₁-C₄ alkylthio, phenoxy, C₁-C₄ haloalkyl, nitro, cyano, C₁-C₄ alkylamino, diloweralkylamino, or C₁-C₄ alkylsulfonyl group or phenyl optionally substituted with one C₁-C₄ alkyl, C₁-C₄ alkoxy or halogen; and when taken together, Y and Z may form a

ring in which YZ are represented by the structure: $-(CH_2)_n-$, where n is an integer of 3 or 4, provided that X is hydrogen; or



where L, M, Q and R are independently hydrogen, halogen, C_1 - C_4 alkyl or C_1 - C_4 alkoxy; and the 2,6-dinitroaniline derivative has the structural formula (II) below



(II)

wherein

- R_3 is hydrogen, C_2 - C_4 alkyl or chloroethyl;
- R_4 is C_2 - C_5 alkyl, chloroethyl, 2-methallyl or cyclopropylmethyl;
- R_5 is hydrogen, methyl or amino;
- R_6 is trifluoromethyl, C_1 - C_3 alkyl, $-SO_2NH_2$ or SO_2CH_3 .

3. The composition according to Claim 2 comprising on a weight basis 2% to 6% of the ethylene oxide/propylene oxide block copolymer, 1% to 6% of the alkylphenol polyethylene oxide condensate, 0.5% to 10% of the antigellation agent and a sufficient amount of the aromatic solvent to total 100%.
4. The composition according to Claim 3 comprising on a weight basis 20% to 40% of the 2,6-dinitroaniline derivative.
5. The composition according to Claim 4 wherein the alkylarylsulfonic acid is a C_8 - C_{18} alkylbenzenesulfonic acid, the ethylene oxide/propylene oxide block copolymer is a butyl-omega-hydroxy-poly(oxypropylene) block polymer with poly(oxyethylene), the alkylphenol polyethylene oxide condensate is a nonylphenol ethoxylate, the antigellation agent is N-methylpyrrolidone, cyclohexanone, an alcohol or a glycol and the aromatic solvent is an aromatic hydrocarbon mixture having a distillation range of from about 140°C to 305°C.
6. The composition according to Claim 5 wherein the C_8 - C_{18} alkylbenzenesulfonic acid is dodecylbenzenesulfonic acid, the butyl-omega-hydroxy-poly(oxypropylene) block polymer with poly(oxyethylene) is alpha-butyl-omega-hydroxy-ethylene oxide-propylene oxide block copolymer having a HLB of 12 and a viscosity at 25°C of 2,000 CPS, the nonylphenol ethoxylate is nonylphenol ethoxylate (9 to 10 mols of ethylene oxide), the antigellation agent is N-methylpyrrolidone, and the aromatic hydrocarbon mixture has a distillation range of from about 183°C to 280°C.
7. The composition according to Claim 5 wherein the imidazoliny acid is selected from the group consisting of 2-(4-isopropyl-4-methyl-5-oxo-2-imidazolin-2-yl)-3-quinolinecarboxylic acid, 5-ethyl-2-(4-isopropyl-4-methyl-5-oxo-2-imidazolin-2-yl)nicotinic acid, 2-(4-isopropyl-4-methyl-5-oxo-2-imidazolin-2-yl)nicotinic acid and 2-(4-isopropyl-4-methyl-5-oxo-2-imidazolin-2-yl)-5-(methoxymethyl) nicotinic acid; and the 2,6-dinitroaniline derivative is selected from the group consisting of pendimethalin and trifluralin.

8. The composition according to Claim 7 wherein the imidazolinyl acid is 5-ethyl-2-(4-isopropyl-4-methyl-5-oxo-2-imidazolin-2-yl)nicotinic acid and the 2,6-dinitroaniline derivative is pendimethalin.

9. The composition according to Claim 7 wherein the imidazolinyl acid is 5-ethyl-2-(4-isopropyl-4-methyl-5-oxo-2-imidazolin-2-yl)nicotinic acid and the 2,6-dinitroaniline derivative is trifluralin.

10. A method of preparing a herbicidal emulsifiable concentrate composition comprising admixing 0.5% to 5% by weight of an imidazolinyl acid; 0% to about 40% by weight of a 2,6-dinitroaniline derivative; 2% to 6% by weight of an alkylarylsulfonic acid; ethylene oxide/propylene oxide block copolymer and alkylphenol polyethylene oxide condensate emulsifying agents; an antigellation agent; and an aromatic solvent.



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number

EP 91 12 1536

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.5)
D,A	US-A-4 822 405 (C.A. MARTIN ET. AL.)		A01N43/50
P,A	EP-A-0 433 577 (AMERICAN CYANAMID COMPANY)		/(A01N43/50,
	---		33:18, 33:20,
A	EP-A-0 249 075 (AMERICAN CYANAMID COMPANY)		25:30, 25:02)

			TECHNICAL FIELDS SEARCHED (Int. Cl.5)
			A01N
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 07 MAY 1992	Examiner DONOVAN T.M.
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone		T : theory or principle underlying the invention	
Y : particularly relevant if combined with another document of the same category		E : earlier patent document, but published on, or after the filing date	
A : technological background		D : document cited in the application	
O : non-written disclosure		L : document cited for other reasons	
P : intermediate document		* : member of the same patent family, corresponding document	